

THE PAPUA NEW GUINEA UNIVERSITY OF TECHNOLOGY

FIRST SEMESTER EXAMINATION – 2022

CH211 – APPLIED PHYSICAL CHEMISTRY

FRIDAY 10th JUNE - 12:50 PM

**TIME ALLOWED: 2 HOURS**

**INFORMATION FOR CANDIDATES:**

1. You will have 10 minutes to read the question paper. You **MUST NOT** begin writing in the answer book during this time.
2. **ANSWER ALL QUESTIONS.**
3. All answers **MUST** be written on the answer book provided.
4. Calculators are permitted in the examination room. Lecture notes, notebooks, plain papers, and textbooks are **NOT** allowed.
5. Mobile phones are not allowed. **SWITCH OFF THE MOBILE PHONES.**
6. Show all working and calculations in the answer book.
7. **DRAW any FIGURES** clearly and visibly.
8. Write your name and number clearly on the front page of the answer book. **DO IT NOW.**

**MARKING SCHEME: [TOTAL: 50 MARKS]**

**Question 1:**

(a) Mention three key features of a Second Order kinetic reaction.

[3 marks]

(b) The following rates of reactions were obtained in FOUR experiments for the reaction  $2A + 2B + C \rightarrow D + 3E$

Experiment	Initial [A] (mol/L)	Initial [B] (mol/L)	Initial [C] (mol/L)	Initial Rate Ms <sup>-1</sup>
1	0.024	0.085	0.032	$6.0 \times 10^{-6}$
2	0.096	0.085	0.032	$9.6 \times 10^{-5}$
3	0.024	0.034	0.080	$1.5 \times 10^{-5}$
4	0.012	0.170	0.032	$1.5 \times 10^{-6}$

(i) Using a simple calculation, deduce the order of the reaction with respect to specie A. Show your deduction very clearly.

[2 marks]

(ii) As in (b) (i) above, deduce the order of the reaction with respect to specie B. Show your deduction very clearly.

[2 marks]

(iii) As in (b) (i) above, deduce the order of the reaction with respect to specie C. Show your deduction very clearly.

[2 marks]

(iv) From your results in (b) (i), (ii), (iii), write the full rate law for the reaction.

[1 mark]

(v) From (b) (iv) above, calculate the rate constant,  $k$ , for the reaction.

[2 marks]

(TOTAL: 12 MARKS)

**GO TO THE NEXT PAGE FOR QUESTION 2**

**Question 2:**

Consider the industrial reaction  $\text{CO(g)} + \text{H}_2\text{O(g)} \leftrightarrow \text{CO}_2\text{(g)} + \text{H}_2\text{(g)}$ . Engineers use the extent of the change from CO and H<sub>2</sub>O to CO<sub>2</sub> and H<sub>2</sub> to regulate the proportions of synthetic fuel mixtures in the industrial reaction.

- (a) If 0.500 mol of CO and 0.500 mol of H<sub>2</sub>O gases are placed in a 250 mL flask at 900 K. When equilibrium is achieved, what concentration of each specie will be present in the reaction vessel? You are given that at 900 K the  $K_c$  for the reaction is 1.56.

[12 marks]

- (b) As a Chemist you proposed that increasing pressure will favour the production of more H<sub>2</sub>(g) as the key fuel of interest. Must a company invest in a high-pressure chemical plant for this industrial process?

[3 marks]

(TOTAL: 15 MARKS)

**Question 3:**

Use the thermodynamic data presented in the table below to calculate  $\Delta S^\circ$

Universe for the reaction:  $\text{PCl}_3\text{(g)} + \text{Cl}_2\text{(g)} \rightarrow \text{PCl}_5\text{(g)}$ . All data are at 298 K.

Show all calculations clearly. [ $\Delta S^\circ(\text{universe}) = \Delta S^\circ(\text{surrounding}) + \Delta S^\circ(\text{system})$ ]

	$\Delta H_f^\circ$ (kJ/mol)	$\Delta G_f^\circ$ (kJ/mol)
PCl <sub>3</sub> (g)	-287	-268
Cl <sub>2</sub> (g)	0	0
PCl <sub>5</sub> (g)	-402	-323

(TOTAL: 5 Marks)

**GO TO THE NEXT PAGE FOR QUESTION 4**

**Question 4:**

Consider the thermodynamic data below and use the data for parts (a), (b) and (c) below:

	$\Delta H_f^\circ$ (kJ/mol)	$\Delta G_f^\circ$ (kJ/mol)
$\text{N}_2\text{O}_4(\text{g})$	9.16	97.89
$\text{O}_2(\text{g})$	0	0
$\text{N}_2\text{O}_5(\text{g})$	11.3	115.1

- (a) Calculate  $\Delta G^\circ$  at 298 K for the reaction,  $2\text{N}_2\text{O}_4(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2\text{N}_2\text{O}_5(\text{g})$ .
- (b) Calculate  $K_p$  at 298 K for the reaction in (a) above. [ $R = 8.314 \text{ J/mol/K}$ ;  
 $\Delta G^\circ = -RT \ln K_p$ ]
- (b) Calculate  $K_c$  for the reaction at 298 K? [ $K_p = K_c(RT)^{\Delta n}$ ]

(TOTAL: 6 MARKS)

**GO TO THE NEXT PAGE FOR QUESTION 5**

**Question 5:**

- (a) You are an Industrial Chemist in charge of a municipal wastewater plant. You have found the wastewater to contain lots of the following ions: Cadmium ( $\text{Cd}^{2+}$ ), Nickel ( $\text{Ni}^{2+}$ ), Manganese ( $\text{Mn}^{2+}$ ), and Calcium ( $\text{Ca}^{2+}$ ). These ions need to be removed by selective precipitation (one ion precipitating at a time) using aqueous NaOH to precipitate them as hydroxides. Therefore, you found the following information presented in the table below.

Chemical Substance	Solubility Product, $K_{sp}$
Calcium hydroxide, $\text{Ca}(\text{OH})_2$	$6.5 \times 10^{-6}$
Cadmium hydroxide, $\text{Cd}(\text{OH})_2$	$7.2 \times 10^{-15}$
Nickel (II) hydroxide, $\text{Ni}(\text{OH})_2$	$6.0 \times 10^{-16}$
Manganese (II) hydroxide, $\text{Mn}(\text{OH})_2$	$1.6 \times 10^{-13}$

- (i) Which of the ions in the wastewater will precipitate first on addition of the aqueous NaOH? Why? [3 marks]
- (ii) Which of the ions will precipitate next with addition of aqueous NaOH? Why? [3 marks]
- (b) Given that the  $K_{sp}$  of  $\text{Ca}_3(\text{PO}_4)_2$  is  $1.2 \times 10^{-29}$  calculate the molar solubility of  $\text{Ca}_3(\text{PO}_4)_2$  in pure water. [6 marks]

(TOTAL: 12 MARKS)

-----THE END. -----