



THE PAPUA NEW GUINEA UNIVERSITY OF TECHNOLOGY
DEPARTMENT OF MATHEMATICS & COMPUTER SCIENCE
SECOND SEMESTER EXAMINATIONS - 2022

FIRST YEAR ELECTRICAL AND MINING ENGINEERING AND MINERAL
PROCESSING

EN121A – ENGINEERING MATHEMATICS II

TIME ALLOWED: 3 HOURS

INFORMATION FOR CANDIDATES

1. Write your name and student number clearly on the front of the examination booklet.
2. You have 10 minutes to read this paper. You must not begin writing during this time.
3. **Answer any four (4) questions out of five (5) questions.**
4. All answers must be written in examination booklets only. No other written material will be accepted.
5. Start the answer for each question on a **new** page. Do **not** use red ink.
6. Notes and textbooks are not allowed in the examination room. All mobile phones and electronic/recording devices must be switched off during the examination.
7. Scientific calculators are allowed in the examination room.
8. A 3 (three) page formula sheet is attached.

MARKING SCHEME


Marks are indicated at the beginning of each question. All questions carry equal marks.

Question 1**MATRICES****(25 marks)**

a) Given,

$$\mathbf{A} = \begin{bmatrix} 1 & 2 & 5 \\ 0 & -1 & 2 \\ 2 & 4 & 11 \end{bmatrix}$$

- i) Calculate the inverse from $\mathbf{A}^{-1} = \frac{1}{\det \mathbf{A}} [A_{jk}]^T$ where A_{jk} is the minor of a_{jk} in $\det \mathbf{A}$. **(8 marks)**
- ii) Check by using $\mathbf{A}\mathbf{A}^{-1} = \mathbf{I}$ (Show all steps). **(7 marks)**
- b) Using Kirchoff's laws, find the currents in the following networks. **(10 marks)**




Question 2

VECTORS

(25 marks)

- a) Find the work done by a force $\mathbf{p} = [2, 6, 6]$ acting on a body if the body is displaced from a point to a point B along the straight segment AB , where $A: (3, 4, 0)$ & $B: (5, 8, 0)$. Sketch \mathbf{p} and AB . Show the details of your work. **(7 marks)**
- b) In the following w.r.t a right-handed Cartesian coordinate system, let $\mathbf{a} = [1, 2, 0]$, $\mathbf{b} = [-3, 2, 0]$. Find the following expressions.
- i) $\mathbf{a} \times \mathbf{b}$, **(4 marks)**
- ii) $\mathbf{a} \cdot \mathbf{b}$ **(4 marks)**
- c)



The access door is held in the 30° open position by the chain AB . If the tension in the chain is 100 N, determine the projection of the tension force on the diagonal axis CD of the door. *(Hint: $F_{CD} = \vec{F}_{AB} \cdot \vec{n}_{CD}$)* **(10 marks)**

Question 3 BASIC LINEAR HOMOGENEOUS 1ST ORDER ODE (25 marks)

- a) Solve $y' = \frac{1-2y-4x}{1+y+2x}$. Hint, use $y + 2x = v$ **(13marks)**
- b) Initially 100mg of radioactive substance was present. After 6 hours the mass had decreased by 3%. If the rate of decay is proportional to the amount of the substance present at time t ,
- i) find the amount remaining after 24 hours. **(7 marks)**
- ii) Determine the half-life of the radioactive substance. **(5 marks)**

Question 4 **1ST AND 2ND ORDER ODE** **(25 marks)**

a) The follow homogeneous equations can be classed as exact differential equations.

$$ye^x dx + (2y + e^x)dy = 0, \quad y(0) = -1$$

As such,

- i) demonstrate exactness, **(4 marks)**
- ii) Find the implicit solution u by appropriate integration and find the constant function, **(4 marks)**
- iii) check your answer by implicit differentiation. **(4 marks)**

b) Find the transient oscillations of the mass-spring system governed by the given equation.

$$y'' + 2y' + 5y = -13 \sin 3t$$

(13 marks)

Question 5 **THE LAPLACE TRANSFORM** **(25 marks)**

Find the Laplace transforms of the following functions (using the Laplace table provided).

- a) $2t + 6$ **(3 marks)**
- b) $\sin 2t \cos 2t$ *Hint: make use of an identity* **(3 marks)**
- c) Given $F(s) = \mathcal{L}(f) = \frac{5s}{s^2 - 25}$, find $f(t)$. **(4 marks)**

In the following system of differential equations, solve the given initial value problems by means of Laplace transforms.

- d) $y_1' = -y_1 + y_2, \quad y_2' = -y_1 - y_2, \quad y_1(0) = 1, \quad y_2(0) = 0$ **(15 marks)**

DATA SHEET for EN121A EXAM 2022 SEMESTER 2

Derivatives and Integrals

<p>1 $\frac{d}{dx}(x^n) = nx^{n-1}$</p> <p>2 $\frac{d}{dx}(\ln x) = \frac{1}{x}$</p> <p>3 $\frac{d}{dx}(e^x) = e^x$</p> <p>4 $\frac{d}{dx}(e^{kx}) = ke^{kx}$</p> <p>5 $\frac{d}{dx}(a^x) = a^x \ln a$</p> <p>6 $\frac{d}{dx}(\cos x) = -\sin x$</p> <p>7 $\frac{d}{dx}(\sin x) = \cos x$</p> <p>8 $\frac{d}{dx}(\tan x) = \sec^2 x$</p> <p>9 $\frac{d}{dx}(\cosh x) = \sinh x$</p> <p>10 $\frac{d}{dx}(\sinh x) = \cosh x$</p> <p>11 $\frac{d}{dx}(\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}$</p> <p>12 $\frac{d}{dx}(\cos^{-1} x) = -\frac{1}{\sqrt{1-x^2}}$</p> <p>13 $\frac{d}{dx}(\tan^{-1} x) = \frac{1}{1+x^2}$</p> <p>14 $\frac{d}{dx}(\sinh^{-1} x) = \frac{1}{\sqrt{x^2+1}}$</p> <p>15 $\frac{d}{dx}(\cosh^{-1} x) = \frac{1}{\sqrt{x^2-1}}$</p> <p>16 $\frac{d}{dx}(\tanh^{-1} x) = \frac{1}{1-x^2}$</p>	<p>$\therefore \int x^n dx = \frac{x^{n+1}}{n+1} + C \quad \left\{ \begin{array}{l} \text{provided} \\ n \neq -1 \end{array} \right\}$</p> <p>$\therefore \int \frac{1}{x} dx = \ln x + C$</p> <p>$\therefore \int e^x dx = e^x + C$</p> <p>$\therefore \int e^{kx} dx = \frac{e^{kx}}{k} + C$</p> <p>$\therefore \int a^x dx = \frac{a^x}{\ln a} + C$</p> <p>$\therefore \int \sin x dx = -\cos x + C$</p> <p>$\therefore \int \cos x dx = \sin x + C$</p> <p>$\therefore \int \sec^2 x dx = \tan x + C$</p> <p>$\therefore \int \sinh x dx = \cosh x + C$</p> <p>$\therefore \int \cosh x dx = \sinh x + C$</p> <p>$\therefore \int \frac{1}{\sqrt{1-x^2}} dx = \sin^{-1} x + C$</p> <p>$\therefore \int \frac{-1}{\sqrt{1-x^2}} dx = \cos^{-1} x + C$</p> <p>$\therefore \int \frac{1}{1+x^2} dx = \tan^{-1} x + C$</p> <p>$\therefore \int \frac{1}{\sqrt{x^2+1}} dx = \sinh^{-1} x + C$</p> <p>$\therefore \int \frac{1}{\sqrt{x^2-1}} dx = \cosh^{-1} x + C$</p> <p>$\therefore \int \frac{1}{1-x^2} dx = \tanh^{-1} x + C$</p>
-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

More derivatives

$$\frac{d}{dx}[\tan x] = \sec^2 x$$

$$\frac{d}{dx}[\sec x] = \sec x \tan x$$

$$\frac{d}{dx}[\cot x] = -\csc^2 x$$

$$\frac{d}{dx}[\csc x] = -\csc x \cot x$$

Specific integrals

$$\int \tan x dx = \ln|\sec x| + C$$

$$\int \sec x dx = \ln|\sec x + \tan x| + C$$

TRIGONOMETRIC SUBSTITUTIONS

EXPRESSION IN THE INTEGRAND	SUBSTITUTION	RESTRICTION ON θ	SIMPLIFICATION
$\sqrt{a^2 - x^2}$	$x = a \sin \theta$	$-\pi/2 \leq \theta \leq \pi/2$	$a^2 - x^2 = a^2 - a^2 \sin^2 \theta = a^2 \cos^2 \theta$
$\sqrt{a^2 + x^2}$	$x = a \tan \theta$	$-\pi/2 < \theta < \pi/2$	$a^2 + x^2 = a^2 + a^2 \tan^2 \theta = a^2 \sec^2 \theta$
$\sqrt{x^2 - a^2}$	$x = a \sec \theta$	$\begin{cases} 0 \leq \theta < \pi/2 & (\text{if } x \geq a) \\ \pi/2 < \theta \leq \pi & (\text{if } x \leq -a) \end{cases}$	$x^2 - a^2 = a^2 \sec^2 \theta - a^2 = a^2 \tan^2 \theta$

Reduction formula

$$\int \sin^n x dx = -\frac{1}{n} \sin^{n-1} x \cos x + \frac{n-1}{n} \int \sin^{n-2} x dx$$

$$\int \cos^n x dx = \frac{1}{n} \cos^{n-1} x \sin x + \frac{n-1}{n} \int \cos^{n-2} x dx$$

$$\int \tan^n x dx = \frac{\tan^{n-1} x}{n-1} - \int \tan^{n-2} x dx$$

$$\int \sec^n x dx = \frac{\sec^{n-2} x \tan x}{n-1} + \frac{n-2}{n-1} \int \sec^{n-2} x dx$$

INTEGRATING PRODUCTS OF TANGENTS AND SECANTS

$\int \tan^m x \sec^n x dx$	PROCEDURE	RELEVANT IDENTITIES
n even	<ul style="list-style-type: none"> Split off a factor of $\sec^2 x$. Apply the relevant identity. Make the substitution $u = \tan x$. 	$\sec^2 x = \tan^2 x + 1$
m odd	<ul style="list-style-type: none"> Split off a factor of $\sec x \tan x$. Apply the relevant identity. Make the substitution $u = \sec x$. 	$\tan^2 x = \sec^2 x - 1$
$\begin{cases} m \text{ even} \\ n \text{ odd} \end{cases}$	<ul style="list-style-type: none"> Use the relevant identities to reduce the integrand to powers of $\sec x$ alone. Then use the reduction formula for powers of $\sec x$. 	$\tan^2 x = \sec^2 x - 1$

Trigonometrical identities

(a) $\sin^2 \theta + \cos^2 \theta = 1$; $\sec^2 \theta = 1 + \tan^2 \theta$; $\operatorname{cosec}^2 \theta = 1 + \cot^2 \theta$

(b) $\sin(A + B) = \sin A \cos B + \cos A \sin B$

$\sin(A - B) = \sin A \cos B - \cos A \sin B$

$\cos(A + B) = \cos A \cos B - \sin A \sin B$

$\cos(A - B) = \cos A \cos B + \sin A \sin B$

$\tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$

$\tan(A - B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$

(c) Let $A = B = \theta \therefore \sin 2\theta = 2 \sin \theta \cos \theta$

$\cos 2\theta = \cos^2 \theta - \sin^2 \theta = 1 - 2 \sin^2 \theta = 2 \cos^2 \theta - 1$

$\tan 2\theta = \frac{2 \tan \theta}{1 - \tan^2 \theta}$

Hyperbolic identities

$\cosh x + \sinh x = e^x$

$\cosh x - \sinh x = e^{-x}$

$\cosh^2 x - \sinh^2 x = 1$

$1 - \tanh^2 x = \operatorname{sech}^2 x$

$\operatorname{coth}^2 x - 1 = \operatorname{csch}^2 x$

$\cosh(-x) = \cosh x$

$\sinh(-x) = -\sinh x$

$\sinh(x + y) = \sinh x \cosh y + \cosh x \sinh y$

$\cosh(x + y) = \cosh x \cosh y + \sinh x \sinh y$

$\sinh(x - y) = \sinh x \cosh y - \cosh x \sinh y$

$\cosh(x - y) = \cosh x \cosh y - \sinh x \sinh y$

$\sinh 2x = 2 \sinh x \cosh x$

$\cosh 2x = \cosh^2 x + \sinh^2 x$

$\cosh 2x = 2 \sinh^2 x + 1 = 2 \cosh^2 x - 1$

- **Kirchhoff's current law:** The algebraic sum of all currents entering and exiting a node must equal zero.
- **Kirchhoff's voltage law:** the voltage around a loop equals the sum of every voltage drop in the same loop for any closed network and equals zero. Ohm's law is: $V = IR$.
- $i \cdot i = j \cdot j = k \cdot k = 1$ & $i \cdot j = j \cdot i = i \cdot k = k \cdot i = j \cdot k = k \cdot j = 0$
- $i \times j = k, j \times k = i, k \times i = j$ & $j \times i = -k, k \times j = -i, i \times k = -j$ & $i \times i = j \times j = k \times k = 0$

• In vector algebra: $\vec{T} = T\vec{n} = T \frac{\overline{AB}}{AB}$

Steps for solving the exact differential equation $u(x, y) = \frac{\partial u}{\partial x} dx + \frac{\partial u}{\partial y} dy$

1. Test for exactness: $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$
2. Find implicit solution $u = \int M dx + k(y)$ or $u = \int N dy + l(x)$. Then find $k'(y)$ or $l'(x)$ with respect to its independent variable alone, and equate to M or N . Then integrate to find value of k or l .
3. Check by implicit differentiation to see if you get the original differential equation $u = M dx + N dy$.

For a non-exact 1st order ODE: $P dx + Q dy = 0$, multiply throughout by integrating factor F to get exact ODE, $FP dx + FQ dy = 0$

F can be calculated as $e^{\int R(x) dx}$ where $R = \frac{1}{Q} \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right)$

For the 1st order non-homogeneous equation, $y' + py = r$,

y is given as $e^{-h} [\int e^{hr} dx + c]$ where $h = \int p dx$

For the 1st order non-linear Bernoulli equation $y' + py = gy^a$ set $u = y^{1-a}$ differentiate this and substitute for y' and y^{1-a} to get as linear nonhomogeneous equation.

Stroud's guide for solving linear 2nd order ode:

- 1 Solution of equations of the form $a \frac{d^2y}{dx^2} + b \frac{dy}{dx} + cy = f(x)$
 - (1) Auxiliary equation: $am^2 + bm + c = 0$
 - (2) Types of solutions:
 - (a) Real and different roots $m = m_1$ and $m = m_2$
 $y = Ae^{m_1x} + Be^{m_2x}$
 - (b) Real and equal roots $m = m_1$ (twice)
 $y = e^{m_1x}(A + Bx)$
 - (c) Complex roots $m = \alpha \pm i\beta$
 $y = e^{\alpha x}(A \cos \beta x + B \sin \beta x)$
- 2 Equations of the form $\frac{d^2y}{dx^2} + n^2y = 0$
 $y = A \cos nx + B \sin nx$
- 3 Equations of the form $\frac{d^2y}{dx^2} - n^2y = 0$
 $y = A \cosh nx + B \sinh nx$
- 4 General solution
 $y = \text{complementary function} + \text{particular integral}$

Kreyszig's guide to solving y_p

Stroud's guide to solving y_p

Table 2.1 Method of Undetermined Coefficients

Term in $r(x)$	Choice for $y_p(x)$
$ke^{\gamma x}$	$Ce^{\gamma x}$
kx^n ($n = 0, 1, \dots$)	$K_n x^n + K_{n-1} x^{n-1} + \dots + K_1 x + K_0$
$k \cos \omega x$	} $K \cos \omega x + M \sin \omega x$
$k \sin \omega x$	
$ke^{\alpha x} \cos \omega x$	} $e^{\alpha x}(K \cos \omega x + M \sin \omega x)$
$ke^{\alpha x} \sin \omega x$	

- | | |
|-----------------------------------|-----------------------------|
| If $f(x) = k \dots \dots$ | Assume $y = C$ |
| $f(x) = kx \dots \dots$ | $y = Cx + D$ |
| $f(x) = kx^2 \dots \dots$ | $y = Cx^2 + Dx + E$ |
| $f(x) = k \sin x$ or $k \cos x$ | $y = C \cos x + D \sin x$ |
| $f(x) = k \sinh x$ or $k \cosh x$ | $y = C \cosh x + D \sinh x$ |
| $f(x) = e^{kx} \dots \dots$ | $y = Ce^{kx}$ |

Brief but required table of Laplace transforms:

	$f(t)$	$\mathcal{L}(f)$		$f(t)$	$\mathcal{L}(f)$
1	1	$1/s$	7	$\cos \omega t$	$\frac{s}{s^2 + \omega^2}$
2	t	$1/s^2$	8	$\sin \omega t$	$\frac{\omega}{s^2 + \omega^2}$
3	t^2	$2!/s^3$	9	$\cosh at$	$\frac{s}{s^2 - a^2}$
4	t^n ($n = 0, 1, \dots$)	$\frac{n!}{s^{n+1}}$	10	$\sinh at$	$\frac{a}{s^2 - a^2}$
5	t^a (a positive)	$\frac{\Gamma(a+1)}{s^{a+1}}$	11	$e^{at} \cos \omega t$	$\frac{s-a}{(s-a)^2 + \omega^2}$
6	e^{at}	$\frac{1}{s-a}$	12	$e^{at} \sin \omega t$	$\frac{\omega}{(s-a)^2 + \omega^2}$

Generally any Laplace transform of a derivative can be written as:

$$\mathcal{L}(f^{(n)}) = s^n \mathcal{L}(f) - s^{n-1} f(0) - s^{n-2} f'(0) - \dots - f^{(n-1)}(0)$$